

EOS Toolkit References:

AAP Endorsement of CDC Guideline. <http://www.aap.org/policy/groupb.html>

ACOG Endorsement of CDC Guideline:

http://www.acog.org/from_home/publications/press_releases/nr11-29-02-1.cfm

Adair CE et al: Risk factors for early-onset group B streptococcal disease in neonates: a population-based case-control study. *CMAJ* 2003;169:198-203.

Aharoni A, Potasman I, Levitan Z, Golan D, Sharf M. Postpartum maternal group B streptococcal meningitis. *Rev Infect Dis* 1990;12:273-6.

Aitmhand R, Moustouli N, Belabbes H, Elmdaghri N, Benbachir M. Serotypes and antimicrobial susceptibility of group B streptococcus isolated from neonates in Casablanca. *Scand J Infect Dis* 2000;32: 339-40.

American College of Obstetricians and Gynecologists, Committee on Obstetric Practice. Prevention of early-onset group B streptococcal disease in newborns [Opinion 173]. Washington, D. C: American College of Obstetricians and Gynecologists, 1996.

American Academy of Pediatrics, Committee on Infectious Diseases/ Committee on Fetus and Newborn. Revised guidelines for prevention of early-onset group B streptococcal (GBS) disease. *Pediatrics* 1997;99:489-96.

Andrews JJ, Diekema DJ, Hunter SK, et al. Group B streptococci causing neonatal bloodstream infection: antimicrobial susceptibility and serotyping results from SENTRY centers in the Western Hemisphere. *Am J Obstet Gynecol* 2000;183:859-62.

Angstetra D, Ferguson J, Giles WB. Institution of universal screening for Group B streptococcus (GBS) from a risk management protocol results in reduction of early-onset GBS disease in a tertiary obstetric unit. *Aust N Z J Obstet Gynaecol* 2007;47:378-82.

Badri MS, Zawaneh S, Cruz AC, et al. Rectal colonization with group B streptococcus: relation to vaginal colonization of pregnant women. *J Infect Dis* 1977;135:308-12.

Baker CJ, Barrett FF, Gordon RC, Yow MD. Suppurative meningitis due to streptococci of Lancefield group B: a study of 33 infants. *J Pediatr* 1973;82:724-9.

- Baker CJ, Edwards MS, Kasper DL. Role of antibody to native type III polysaccharide of group B streptococcus in infant infection. *Pediatrics* 1981;68:544–9.
- Baker CJ, Edwards MS. Group B streptococcal infections. In: Remington J, Klein JO, eds. *Infectious diseases of the fetus and newborn infant*. Philadelphia: W.B. Saunders, 1990:742–811.
- Baker CJ, Paoletti LC, Rench MA, et al. Use of capsular polysaccharide– tetanus toxoid conjugate vaccine for type II group B streptococcus in healthy women. *J Infect Dis* 2000;182:1129–38.
- Baker CJ, Rench MA, Edwards MS, Carpenter RJ, Hays BM, Kasper DL. Immunization of pregnant women with a polysaccharide vaccine of group B streptococcus. *N Engl J Med* 1988;319:1180–5.
- Baker CJ, Paoletti LC, Wessels MR, et al. Safety and immunogenicity of capsular polysaccharide–tetanus toxoid conjugate vaccines for group B streptococcal types Ia and Ib. *J Infect Dis* 1999;179:142–50.
- Baltimore, R. S.; Huie, S. M.; Meek, J. I.; Schuchat, A., and O'Brien, K. L. Early-onset neonatal sepsis in the era of group B streptococcal prevention. *Pediatrics*. 2001 Nov; 108(5):1094-8.
- Balter et al: Impact of intrapartum antibiotics on the care and evaluation of the neonate. *Pediatr Infect Dis J* 2003; 22:853-7.
- Balter S, Zell E, O'Brien K, et al. Evaluating the impact of intrapartum antibiotics to prevent group B streptococcus on the care and workup of the neonate [Abstract]. In: *Program and Abstracts of the 40th Interscience Conference on Antimicrobial Agents and Chemotherapy*. Washington DC: American Society for Microbiology, 2000:460.
- Benitz WE, Gould JB, Druzin ML. Antimicrobial prevention of earlyonset group B streptococcal sepsis: estimates of risk reduction based on a critical literature review. *Pediatrics* 1999;103:e78. Available at <http://www.pediatrics.org/cgi/content/full/103/6/e78>.
- Bergeron MG, Ke D, Menard C, et al. Rapid detection of group B streptococci in pregnant women at delivery. *N Engl J Med* 2000;343:175–9.
- Berkowitz K, Regan JA, Greenberg E. Antibiotic resistance patterns of group B streptococci in pregnant women. *J Clin Microbiol* 1990;28:5–7.

- Bland et al: Antibiotic resistance patterns of Group B Streptococci in late third-trimester rectovaginal cultures. *Am J Obstet Gynecol*. 2001;184:1125-6.
- Bobitt JR, Ledger WJ. Amniotic fluid analysis: its role in maternal and neonatal infection. *Obstet Gynecol* 1978;51:56–62.
- Boulvain M, Stan C, Irion O. Membrane sweeping for induction of labour (Cochrane Review). In: *The Cochrane Library*, Issue 4. Oxford: 2001.
- Boyer KM, Gadzala CA, Kelly PD, Burd LI, Gotoff SP. Selective intrapartum chemoprophylaxis of neonatal group B streptococcal earlyonset disease. II. Predictive value of prenatal cultures. *J Infect Dis* 1983;148:802–9.
- Boyer KM, Gadzala CA, Burd LI, Fisher DE, Paton JB, Gotoff SP. Selective intrapartum chemoprophylaxis of neonatal group B streptococcal early-onset disease. I. Epidemiologic rationale. *J Infect Dis* 1983;148:795–801.
- Boyer KM, Gotoff SP. Strategies for chemoprophylaxis of GBS earlyonset infections. *Antibiot Chemother* 1985;35:267–80.
- Boyer KM, Gotoff SP. Prevention of early-onset neonatal group B streptococcal disease with selective intrapartum chemoprophylaxis. *N Engl J Med* 1986;314:1665–9.
- Braun TI, Pinover W, Sih P. Group B streptococcal meningitis in a pregnant woman before the onset of labor. *Clin Infect Dis* 1995;21:1042–3.
- Bromberger P, Lawrence JM, Braun D, Saunders B, Contreras R, Petitti DB. The influence of intrapartum antibiotics on the clinical spectrum of early-onset group B streptococcal infection in term infants. *Pediatrics* 2000;106:244–50.
- Broad-spectrum antibiotics for spontaneous preterm labour: the ORACLE II randomised trial. ORACLE Collaborative Group. *Lancet*. 2001 Mar 31; 357(9261):989-94.
- Castor ML, Whitney C, Facklam R, et al. Antimicrobial susceptibility and serotype patterns of invasive group B Streptococcus isolates from Georgia, Minnesota, New York and Oregon, 1996–2000 [Abstract]. *International Conference on Emerging Infectious Diseases 2002*, program and abstract book. Atlanta: 2002, 132.
- CDC. Adoption of hospital policies for prevention of perinatal group B streptococcal disease—United States, 1997. *MMWR* 1998;47: 665–70.
- CDC. Adoption of perinatal group B streptococcal disease prevention recommendations by prenatal-care providers—Connecticut and Minnesota, 1998. *MMWR* 2000;49:228–32.

- CDC. Early-onset group B streptococcal disease, United States, 1998– 1999. *MMWR* 2000;49:793–6.
- CDC. Hospital-based policies for prevention of perinatal group B streptococcal disease—United States, 1999. *MMWR* 2000;49:936–40.
- CDC. Laboratory practices for prenatal group B streptococcal screening and reporting—Connecticut, Georgia, and Minnesota, 1997–1998. *MMWR* 1999;48:426–8.
- CDC. Prevention of perinatal group B streptococcal disease: a public health perspective. *MMWR* 1996;45 (RR-7):1–24.
- CDC. Sexually transmitted diseases treatment guidelines, 2002. *MMWR* 2002;51 (No. RR- 6):28–9.
- Chen et al: The role of intrapartum fever in identifying asymptomatic term neonates with early- onset neonatal sepsis. *J Perinatol.* 2002;22:653-7.
- Chen KT, Puopolo KM, Eichenwald EC, Onderdonk AB, Lieberman E. No increase in rates of early-onset neonatal sepsis by antibiotic-resistant group B *Streptococcus* in the era of intrapartum antibiotic prophylaxis. *Am J Obstet Gynecol* 2005;192:1167-71.
- Chen, K. T.; Tuomala, R. E.; Cohen, A. P.; Eichenwald, E. C., and Lieberman, E. No increase in rates of early-onset neonatal sepsis by non-group B *Streptococcus* or ampicillin-resistant organisms. *Am J Obstet Gynecol.* 2001 Oct; 185(4):854-8.
- Cheon-Lee E, Amstey MS. Compliance with the Centers for Disease Control and Prevention antenatal culture protocol for preventing group B streptococcal neonatal sepsis. *Am J Obstet Gynecol* 1998;179:77–9.
- Chow, L. C.; Wright, K. W., and Sola, A. Can changes in clinical practice decrease the incidence of severe retinopathy of prematurity in very low birth weight infants? *Pediatrics.* 2003 Feb; 111(2):339-45.
- Christensen KK, Dahlander K, Linden V, Svenningsen N, Christensen P. Obstetrical care in future pregnancies after fetal loss in group B streptococcal septicemia. A prevention program based on bacteriological and immunological follow-up. *Eur J Obstet Gynecol Reprod Biol* 1981;12:143–50.
- Clemens et al: The development of a group B streptococcus prevention policy at a community hospital. *J Perinatol.* 2002; 22:523-5.

- Cordero L, Sananes M, Ayers LW. Bloodstream infections in a neonatal intensive-care unit: 12 years' experience with an antibiotic control program. *Infect Control Hosp Epidemiol* 1999;20:242–6.
- Daily P; Gelling L, and Mukerjee N. Adoption of Hospital Policies for Prevention of Perinatal Group B streptococcal Disease,-United States, 1997. *Morbidity and Mortality Weekly Report*. 1998 Aug 21; 47(32):666-670.
- Davies HD, Adair C, McGeer A, et al. Antibodies to capsular polysaccharides of group B Streptococcus in pregnant Canadian women: relationship to colonization status and infection in the neonate. *J Infect Dis* 2001;184:285–91.
- Davies HD, Adair CE, Schuchat A, Low DE, Sauve RS, McGeer A. Physicians' prevention practices and incidence of neonatal group B streptococcal disease in 2 Canadian regions. *CMAJ* 2001;164: 479–85.
- Davis RL, Hasselquist MB, Cardenas V, et al. Introduction of the new Centers for Disease Control and Prevention group B streptococcal prevention guideline at a large West Coast health maintenance organization. *Am J Obstet Gynecol* 2001;184:603–10. 2001;164: 479–85.
- de Cueto M, Sanchez M-J, Sampedro A, Miranda J-A, Herruzo A-J, Rosa-Fraile M. Timing of intrapartum ampicillin and prevention of vertical transmission of group B streptococcus. *Obstet Gynecol* 1998;91:112–4.
- Dunn AB, Blomquist J, Khouzami V. Anaphylaxis in labor secondary to prophylaxis against group B streptococcus: a case report. *J Reprod Med* 1999;44:381–4.
- Early-onset and late-onset neonatal group B streptococcal disease--United States, 1996-2004. *MMWR Morb Mortal Wkly Rep* 2005;54:1205-8.
- Easmon CS, Hastings MJ, Deeley J, Bloxham B, Rivers RP, Marwood R. The effect of intrapartum chemoprophylaxis on the vertical transmission of group B streptococci. *Br J Obstet Gynaecol* 1983;90:633–5. Vol. 51 / RR-11
Recommendations and Reports
- Edwards et al: Penicillin has no advantage over ampicillin for use in prophylaxis against early- onset neonatal sepsis. *Am J Obstet Gynecol* 2002; 187:224
- Edwards, R. K.; Clark, P., and Duff, P. Intrapartum antibiotic prophylaxis 2: positive predictive value of antenatal group B streptococci cultures and antibiotic susceptibility of clinical isolates. *Obstet Gynecol*. 2002 Sep; 100(3):540-4.
- Escobar GJ, Li D, Armstrong MA, et al. Neonatal sepsis workups in infants >2000 grams at birth: a population-based study. *Pediatrics* 2000;106:256–63.

- Factor SH, Levine OS, Nassar A, et al. Impact of a risk-based prevention policy on neonatal group B streptococcal disease. *Am J Obstet Gynecol* 1998;179:1568–71.
- Factor SH, Whitney CG, Zywicki S, Schuchat A, the Active Bacterial Core Surveillance Team. Effects of hospital policies based on 1996 group B streptococcal disease consensus guidelines. *Obstet Gynecol* 2000;95:377–82.
- Fargason CA, Peralta-Carcelen M, Rouse DJ, Cutter GR, Goldenberg RL. The pediatric costs of strategies for minimizing the risk of early-onset group B streptococcal disease. *Obstet Gynecol* 1997;90: 347–52.
- Faxelius G, Bremme K, Kvist-Christensen K, Christensen P, Ringertz S. Neonatal septicemia due to group B streptococci—perinatal risk factors and outcome of subsequent pregnancies. *J Perinat Med* 1988;16:423–30.
- Fernandez M, Hickman ME, Baker CJ. Antimicrobial susceptibilities of group B streptococci isolated between 1992 and 1996 from patients with bacteremia or meningitis. *Antimicrob Agents Chemother* 1998;42:1517–9.
- Fox BC. Delayed-onset postpartum meningitis due to group B streptococcus [letter]. *Clin Infect Dis* 1994;19:350.
- Franciosi RA, Knostman JD, Zimmerman RA. Group B streptococcal neonatal and infant infections. *J Pediatr* 1973;82:707–18.
- Garland SM, Fliegner JR. Group B streptococcus (GBS) and neonatal infections: the case for intrapartum chemoprophylaxis. *Aust NZ J Obstet Gynaecol* 1991;31:119–22.
- Gervasio, C. T.; Mantaring, B. S.; Alankar, S., and Shankaran, S. Early-onset neonatal group B streptococcal sepsis: intrapartum antibiotic prophylaxis in the clinical setting. *J Perinatol*. 2001 Jan-2001 Feb 28; 21(1):9-14.
- Gibbs RS, Jones PM, Wilder CJY. Internal fetal monitoring and maternal infection following cesarean section: a prospective study. *Obstet Gynecol* 1978;52:193–7.
- Gilbert et al: Compliance with protocols for prevention of neonatal group B streptococcal sepsis- practicalities and limitations. *Infect Dis Obstet Gynecol*. 2003; 11:1-9.
- Gil EG, Rodriguez MC, Bartolome R, Berjano B, Cabero L, Andreu A. Evaluation of the Granada agar plate for detection of vaginal and rectal group B streptococci in pregnant women. *J Clin Microbiol* 1999;37:2648–51. August 16, 2002

- Gilson GJ, Christensen F, Bekes K, Silva L, Qualls CR. Prevention of group B streptococcus early-onset neonatal sepsis: comparison of the Centers for Disease Control and Prevention screening-based protocol to a risk-based protocol in infants at greater than 37 weeks' gestation. *J Perinatol* 2000;20:491–5.
- Goodman LS, Gilman A, Gilman AG. Goodman and Gilman's *The pharmacologic basis of therapeutics*, Tenth edition. New York: Pergamon Press, 1990:1825. August 16, 2002
- Gotoff SP, Boyer KM. Prevention of early-onset neonatal group B streptococcal disease. *Pediatrics* 1997;99:866–9.
- Gupta K, Scholes D, Stamm WE. Increasing prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in women. *JAMA* 1999;282:325–6.
- Hammerschlag MR, Baker CJ, Alpert S, et al. Colonization with group B streptococci in girls under 16 years of age. *Pediatrics* 1977;60:473–6.
- Hafner E, Sterniste W, Rosen A, et al. Group B streptococci during pregnancy: a comparison of two screening and treatment protocols. *Am J Obstet Gynecol* 1998;179:677–81.
- Hager WD, Schuchat A, Gibbs R, Sweet R, Mead P, Larsen JW. Prevention of perinatal group B streptococcal infection: current controversies. *Obstet Gynecol* 2000;96:141–5.
- Hibbard JU, Shashoua A, Adamczyk C, Ismail M. Cervical ripening with prostaglandin gel and hygroscopic dilators. *Infect Dis Obstet Gynecol* 1998;6:18–24.
- Hristevea L, Booy R, Bowler I, Wilkinson AR. Prospective surveillance of neonatal meningitis. *Arch Dis Child* 1993;69:14–8.
- Hyde TB, Hilger TM, Reingold A, Farley MM, O'Brien KL, Schuchat A. Trends in incidence and antimicrobial resistance of early-onset sepsis: population-based surveillance in San Francisco and Atlanta. *Pediatrics* 2002;110:690–5.
- Isaacs D, Royle JA, Australasian Study Group for Neonatal Infections. Intrapartum antibiotics and early onset neonatal sepsis caused by group B *Streptococcus* and by other organisms in Australia. *Pediatr Infect Dis J* 1999;18:524–8.

Jeffery HE, Lahra MM. Eight-year outcome of universal screening and intrapartum antibiotics for maternal group B streptococcal carriers. *Pediatrics* 1998;101:(1). Available at <http://www.pediatrics.org/cgi/content/full/101/1/e2>. 19

Joseph TA, Pyati SP, Jacobs N. Neonatal early-onset *Escherichia coli* disease: the effect of intrapartum ampicillin. *Arch Pediatr Adolesc Med* 1998;152:35–40. Vol. 51 / RR-11 Recommendations and Reports.

Kasper DL, Paoletti LC, Wessels MR, et al. Immune response to type III group B streptococcal polysaccharide–tetanus toxoid conjugate vaccine. *J Clin Invest* 1996;98:2308–14.22 MMWR

Katz VL, Moos M-K, Cefalo RC, Thorp JM, Bowes WA, Wells SD. Group B streptococci: results of a protocol of antepartum screening and intrapartum treatment. *Am J Obstet Gynecol* 1994;170:521–6.

Kaye, D. Gram-negative sepsis in neonates may be rising since start of GBS prophylaxis. *Clin Infect Dis*. 2001 May 15; 32(10):i-ii.

Kelkar PS, Li JT. Cephalosporin allergy. *N Engl J Med* 2001;345:804–9.

Kenyon SL, Taylor DJ, Tarnow-Mordi W, ORACLE Collaborative Group. Broad spectrum antibiotics for preterm, prelabour rupture of fetal membranes: the ORACLE I randomised trial. *Lancet* 2001;357:979–88.

Kenyon SL, Taylor DJ, Tarnow-Mordi W, ORACLE Collaborative Group. Broad-spectrum antibiotics for spontaneous preterm labour: the ORACLE II randomised trial. *Lancet* 2001;357:989–94.

Lansky, A.; Jones, J. L.; Frey, R. L., and Lindegren, M. L. Trends in HIV testing among pregnant women: United States, 1994-1999. *Am J Public Health*. 2001 Aug; 91(8):1291-3.

Lavin BS. Antibiotic cycling and marketing into the 21st century: a perspective from the pharmaceutical industry. *Infect Control Hosp Epidemiol* 2000;21:S32–5.

Levine, E. M.; Ghai, V.; Barton, J. J., and Strom, C. M. Intrapartum antibiotic prophylaxis increases the incidence of gram-negative neonatal sepsis. *Infect Dis Obstet Gynecol*. 1999; 7(4):210-3.

Levine EM, Ghai V, Barton JJ, Strom CM. Intrapartum antibiotic prophylaxis increases the incidence of gram-negative neonatal sepsis. *Infect Dis Obstet Gynecol* 1999;7:210–13.

- Lieu TA, Mohle-Boetani JC, Ray GT, Ackerson LM, Walton DL. Neonatal group B streptococcal infection in a managed care population. *Obstet Gynecol* 1998;92:21–7.
- Lin FYC, Azimi PH, Weisman LE, et al. Antibiotic susceptibility profiles for group B streptococci isolated from neonates, 1995–1998. *Clin Infect Dis* 2000;31:76–9.
- Lin FYC, Brenner RA, Johnson YR, et al. The effectiveness of riskbased intrapartum chemoprophylaxis for the prevention of early-onset neonatal group B streptococcal disease. *Am J Obstet Gynecol* 2001;184:1204–10.
- Lin F-Y, Philips JB, III, Azimi PH, et al. Level of maternal antibody required to protect neonates against early-onset disease caused by group B Streptococcus type Ia: a multicenter, seroepidemiology study. *J Infect Dis* 2001;184:1022–8.
- Locksmith GJ, Clark P, Duff P. Maternal and neonatal infection rates with three different protocols for prevention of group B streptococcal disease. *Am J Obstet Gynecol* 1999;180:416–22.
- Luck S et al: Estimated early-onset group B streptococcal neonatal disease. *Lancet* 2003;361:1953-4.
- Main, E. K. and Slagle, T. Prevention of early-onset invasive neonatal group B streptococcal disease in a private hospital setting: the superiority of culture-based protocols. *Am J Obstet Gynecol*. 2000 Jun; 182(6):1344-54.
- Matorras R, Garcia-Perea A, OmeZaca F, Diez-Enciso M, Madero R, Usandizaga JA. Intrapartum chemoprophylaxis of early-onset group B streptococcal disease. *Eur J Obstet Gynecol Reprod Biol* 1991;40: 57–62.
- McCracken GH. Group B streptococci: the new challenge in neonatal infections. *J Pediatr* 1973;82:703–6. August 16, 2002
- Mercer BM, Carr TL, Beazley DD, Crouse DT, Sibai BM. Antibiotic use in pregnancy and drug-resistant infant sepsis. *Am J Obstet Gynecol* 1999;181:816–21.
- Mercer BM, Ramsey RD, Sibai BM. Prenatal screening for group B streptococcus. II. Impact of antepartum screening and prophylaxis on neonatal care. *Am J Obstet Gynecol* 1995;173:842–6.
- Mercer BM, Taylor MC, Fricke JL, Baselski VS, Sibai BM. The accuracy and patient preference for self-collected group B streptococcus cultures. *Am J Obstet Gynecol* 1995;173:1325–8.

- Mitchell TF, Pearlman MD, Chapman RL, Bhatt-Mehta V, Faix RG. Maternal and transplacental pharmacokinetics of cefazolin. *Obstet Gynecol* 2001;98:1075–9.
- Mohle-Boetani JC, Schuchat A, Plikaytis BD, Smith JD, Broome CV. Comparison of prevention strategies for neonatal Group B streptococcal infection: a population-based economic approach. *JAMA* 1993;270:1442–8.
- Mohle-Boetani JC, Lieu TA, Ray GT, Escobar G. Preventing neonatal group B streptococcal disease: cost-effectiveness in a health maintenance organization and the impact of delayed hospital discharge for newborns who received intrapartum antibiotics. *Pediatrics* 1999;103:703–10.
- Moore et al: Effects of intrapartum antimicrobial prophylaxis for prevention of group-B-streptococcal disease on the incidence and ecology of early-onset sepsis. *Lancet Infect Dis.* 2003; 3:201-13.
- Morales WJ, Dickey SS, Bornick P, Lim DV. Change in antibiotic resistance of group B streptococcus: impact on intrapartum management. *Am J Obstet Gynecol* 1999;181:310–4.
- NCCLS. Performance Standard for Antimicrobial Susceptibility Testing, M100-S12. Table 2H. Wayne, PA, USA: NCCLS, 2002.
- Neumantis-Keller et al: Limitations of the obstetric group B Streptococcus protocol. *J Reprod Med* 2003; 48:107-11.
- Newton ER, Prihoda TJ, Gibbs RS. Logistic regression analysis of risk factors for intra-amniotic infection. *Obstet Gynecol* 1989;73:571–5.
- Ottolini et al: Utility of complete blood count and blood culture screening to diagnose neonatal sepsis in the asymptomatic at risk newborn. *Ped Infect Dis J* 2003; 22:430-4.
- Pass MA, Gray BM, Dillon HC. Puerperal and perinatal infections with group B streptococci. *Am J Obstet Gynecol* 1982;143:147–52.
- Patel, D. M.; Leblanc, M. H.; Morrison, J. C.; Graves, G. R.; Glick, C. G.; Martin, J. N. Jr; Rhodes, P. G., and Chauhan, S. P. Postnatal penicillin prophylaxis and the incidence of group B streptococcal sepsis in neonates. *South Med J.* 1994 Nov; 87(11):1117-20.
- Patel, D. M.; Rhodes, P. G.; LeBlanc, M. H.; Graves, G. R.; Glick, C., and Morrison, J. Role of postnatal penicillin prophylaxis in prevention of neonatal group B streptococcus infection. *Acta Paediatr.* 1999 Aug; 88(8):874-9.

- Pearlman MD, Pierson CL, Faix RG. Frequent resistance of clinical group B streptococci isolates to clindamycin and erythromycin. *Obstet Gynecol* 1998;92:258–61.
- Peralta-Carcelen M, Fargasan CA, Jr, Cliver SP, Cutter GR, Gigante J, Goldenberg RL. Impact of maternal group B streptococcal screening on pediatric management in full-term newborns. *Arch Pediatr Adolesc Med* 1996;150:802–8.
- Perinatal group B streptococcal disease after universal screening recommendations--United States, 2003-2005. *MMWR Morb Mortal Wkly Rep* 2007;56:701-5.
- Persson K, Christensen KK, Christensen P, Forsgren A, Jorgensen C, Persson PH. Asymptomatic bacteriuria during pregnancy with special reference to group B streptococci. *Scand J Infect Dis* 1985;17:195–9.
- Petrova et al: Obstetrician preferences for prenatal strategies to reduce early-onset group B streptococcal infection in neonates-a population-based survey. *Am J Obstet Gynecol.* 2002; 187:709-14.
- Phares CR, Lynfield R, Farley MM, et al. Epidemiology of Invasive Group B Streptococcal Disease in the United States, 1999-2005. *JAMA* 2008;299:2056-65.
- Pinto, N. M.; Soskolne, E. I.; Pearlman, M. D., and Faix, R. G. Neonatal early-onset group B streptococcal disease in the era of intrapartum chemoprophylaxis: residual problems. *J Perinatol.* 2003 Jun; 23(4):265-71.
- Plsek, P. E. and Greenhalgh, T. Complexity science: The challenge of complexity in health care. *BMJ.* 2001 Sep 15; 323(7313):625-8.
- Puopolo KM, Madoff LC, Eichenwald EC. Early-onset group B streptococcal disease in the era of maternal screening. *Pediatrics* 2005;115:1240-6.
- Pylipow M, Gaddis M, Kinney JS. Selective intrapartum prophylaxis for group B streptococcus colonization: management and outcome of newborns. *Pediatrics* 1994;93:631–5.
- Ramus RM, McIntire DD, Wendel GD, Jr. Antibiotic chemoprophylaxis for group B strep is not necessary with elective cesarean section at term [Abstract]. *Am J Obstet Gynecol* 1999;180:S85.
- Regan JA, Klebanoff MA, Nugent RP, Vaginal Infections and Prematurity Study Group. The epidemiology of group B streptococcal colonization in pregnancy. *Obstet Gynecol* 1991;77:604–10.

- Regan JA, Klebanoff MA, Nugent RP, et al. Colonization with group B streptococci in pregnancy and adverse outcome. *Am J Obstet Gynecol* 1996;174:1354–60.
- Reisner DP, Haas MJ, Zingheim RW, Williams MA, Luthy D. Performance of a group B streptococcal prophylaxis protocol combining high-risk treatment and low-risk screening. *Am J Obstet Gynecol* 2000;182:1335–43.
- Rentz AC, Samore MH, Stoddard GJ, Faix RG, Byington CL. Risk factors associated with ampicillin-resistant infection in newborns in the era of group B streptococcal prophylaxis. *Arch Pediatr Adolesc Med* 2004;158:556–60.
- Riley L, Apollon M, Haider S, et al. “Real world” compliance with strategies to prevent early-onset GBS. Presented at the Infectious Diseases Society for Obstetrics and Gynecology Annual Scientific Meeting. Aug. 9–11, 2001, Quebec City, Canada.
- Riley et al: “Real World” compliance with strategies to prevent early-onset group B streptococcal disease. *J Perinatol*. 2003; 23: 272-7.
- Robbins JB, Schneerson R, Vann WF, Bryla DA, Fattom A. Prevention of systemic infections caused by group B streptococcus and *Staphylococcus aureus* by multivalent polysaccharide–protein conjugate vaccines. *Ann NY Acad Sci* 1995;754:68–82.
- Rosa-Fraile M, Rodriguez-Granger J, Cueto-Lopez M, et al. Use of Granada medium to detect group B streptococcal colonization in pregnant women. *J Clin Microbiol* 1999;37:2674–7.
- Rosenstein NE, Schuchat A. Opportunities for prevention of perinatal group B streptococcal disease: a multistate surveillance analysis. *Obstet Gynecol* 1997;90:901–6.
- Rouse DJ, Goldenberg RL, Cliver SP, Cutter GR, Mennemeyer ST, Fargason CA, Jr. Strategies for the prevention of early-onset neonatal group B streptococcal sepsis: a decision analysis. *Obstet Gynecol* 1994;83:483–94.
- Schrag SJ, Arnold KE, Roome A, et al. Intrapartum antibiotic exposure in the era of perinatal group B streptococcal disease prevention [Abstract G-1824]. In: Program and abstracts of the 41st Annual Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 2001.
- Schrag, S.; Gorwitz, R.; Fultz-Butts, K., and Schuchat, A. Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. *MMWR Recomm Rep*. 2002 Aug 16; 51(RR-11):1-22.

- Schrag SJ, Hadler JL, Arnold KE, Martell-Cleary P, Reingold A, Schuchat A. Risk factors for invasive, early-onset *Escherichia coli* infections in the era of widespread intrapartum antibiotic use. *Pediatrics* 2006;118:570-6.
- Schrag SJ, Whitney CG, Schuchat A. Neonatal group B streptococcal disease: how infection control teams can contribute to prevention efforts. *Infect Control Hosp Epidemiol* 2000;21:473-83. Vol. 51 / RR-11 Recommendations and Reports 23 and 24 MMWR August 16, 2002
- Schrag SJ, Zywicki S, Farley MM, et al. Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. *N Engl J Med* 2000;342:15-20.
- Schrag, S. J.; Zell, E. R.; Lynfield, R.; Roome, A.; Arnold, K. E.; Craig, A. S.; Harrison, L. H.; Reingold, A.; Stefonek, K.; Smith, G.; Gamble, M., and Schuchat, A. A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. *N Engl J Med*. 2002 Jul 25; 347(4):233-9.
- Schuchat A, Deaver-Robinson K, Plikaytis BD, Zangwill KM, Mohle-Boetani J, Wenger JD. Multistate case-control study of maternal risk factors for neonatal group B streptococcal disease. *Pediatr Infect Dis J* 1994;13:623-9.
- Schuchat A. Epidemiology of group B streptococcal disease in the United States: shifting paradigms. *Clin Microbiol Rev* 1998;11: 497-513.
- Schuchat A. Group B streptococcus. *Lancet* 1999;353:51-6.
- Schuchat A, Hilger T, Zell E, et al. Active Bacterial Core Surveillance of the Emerging Infections Program Network. *Emerg Infect Dis* 2001;7:92-9.
- Schuchat A, Oxtoby M, Cochi S, et al. Population-based risk factors for neonatal group B streptococcal disease: results of a cohort study in metropolitan Atlanta. *J Infect Dis* 1990;162:672-7.
- Schuchat A, Roome A, Zell E, Linardos H, Zywicki S, O'Brien KL. Integrated monitoring of a new group B streptococcal disease prevention program and other perinatal infections. *Matern Child Health J* 2002;6:107-14.
- Schuchat A, Wenger JD. Epidemiology of group B streptococcal disease: risk factors, prevention strategies and vaccine development. *Epidemiol Rev* 1994;16:374-402.
- Schuchat A, Zywicki S, Dinsmoor MJ, et al. Risk factors and opportunities for prevention of early-onset neonatal sepsis: a multicenter casecontrol study. *Pediatrics* 2000;105:21-6.

- Siegel JD, Cushion NB. Prevention of early-onset group B streptococcal disease: another look at single-dose penicillin at birth. *Obstet Gynecol* 1996;87:692–8.
- Siegel, J. D.; McCracken, G. H. Jr; Threlkeld, N.; DePasse, B. M., and Rosenfeld, C. R. Single-dose penicillin prophylaxis of neonatal group-B-streptococcal disease. *Lancet*. 1982 Jun 26; 1(8287):1426-30.
- Siegel, J. D.; McCracken, G. H. Jr; Threlkeld, N.; Milvenan, B., and Rosenfeld, C. R. Single-dose penicillin prophylaxis against neonatal group B streptococcal infections. A controlled trial in 18,738 newborn infants. *N Engl J Med*. 1980 Oct 2; 303(14):769-75.
- Silverman NS, Morgan M, Nichols WS. Antibiotic resistance patterns of group B streptococcus in antenatal genital cultures. *J Reprod Med* 2000;45:979–82.
- Sinha et al: Intrapartum antibiotics and neonatal invasive infections caused by organisms other than group B streptococcus. *J Pediatr*. 2003; 142:492-7.
- Soper DE, Mayhall CG, Froggatt JW. Characterization and control of intraamniotic infection in an urban teaching hospital. *Am J Obstet Gynecol* 1996;175:304–10.
- Spaetgens, R.; DeBella, K.; Ma, D.; Robertson, S.; Mucenski, M., and Davies, H. D. Perinatal antibiotic usage and changes in colonization and resistance rates of group B streptococcus and other pathogens. *Obstet Gynecol*. 2002 Sep; 100(3):525-33.
- Stoll, B. J.; Hansen, N.; Fanaroff, A. A.; Wright, L. L.; Carlo, W. A.; Ehrenkranz, R. A.; Lemons, J. A.; Donovan, E. F.; Stark, A. R.; Tyson, J. E.; Oh, W.; Bauer, C. R.; Korones, S. B.; Shankaran, S.; Laptook, A. R.; Stevenson, D. K.; Papile, L. A., and Poole, W. K. Changes in pathogens causing early-onset sepsis in very-low-birth-weight infants. *N Engl J Med*. 2002 Jul 25; 347(4):240-7.
- Stratton KR, Durch JS, Lawrence RS. *Vaccines for the 21st Century: A tool for decisionmaking*. Washington, DC: Institute of Medicine, National Academy Press, 1999.
- Terrone DA, Rinehart BK, Einstein MH, Britt LB, Martin JN, Perry KG. Neonatal sepsis and death caused by resistant *Escherichia coli*: possible consequences of extended maternal ampicillin administration. *Am J Obstet Gynecol* 1999;180:1345–8.
- Towers CV, Carr MH, Padilla G, Asrat T. Potential consequences of widespread antepartal use of ampicillin. *Am J Obstet Gynecol* 1998;179:879–83.

- Towers et al: Antepartum use of antibiotics and early-onset sepsis-the next 4 years. *Am J Obstet Gynecol.* 2002; 187:495-500.
- Towers, C. V. and Briggs, G. G. Antepartum use of antibiotics and early-onset neonatal sepsis: the next 4 years. *Am J Obstet Gynecol.* 2002 Aug; 187(2):495-500.
- Towers, C. V.; Carr, M. H.; Padilla, G., and Asrat, T. Potential consequences of widespread antepartal use of ampicillin. *Am J Obstet Gynecol.* 1998 Oct; 179(4):879-83.
- U. S. Department of Health and Human Services. *Healthy People 2010: understanding and improving health.* 2nd ed. Washington, DC: US Government Printing Office, 2000. Available at [http:// www.health.gov/healthypeople](http://www.health.gov/healthypeople).
- Velaphi et al: Early-onset group B streptococcal infection after a combined maternal and neonatal group B streptococcal chemoprophylaxis strategy. *Pediatrics.* 2003; 111:541-7.
- Visser VE, Hall RT. Lumbar puncture in the evaluation of suspected neonatal sepsis. *J Pediatr* 1980;96:1063-7.
- Votava M, Tejkalová M, Drábková M, Unzeitig V, Braveny I. Use of GBS media for rapid detection of group B streptococci in vaginal and rectal swabs from women in labor. *Eur J Clin Microbiol Infect Dis* 2001;20:120-2.
- Walker CK, Crombleholme WR, Ohm-Smith MJ, Sweet RL. Comparison of rapid tests for detection of group B streptococcal colonization. *Am J Perinatol* 1992;9:304-8.
- Watt JP, Schuchat A, Erickson K, Honig JE, Gibbs R, Schulkin J. Group B streptococcal disease prevention practices of obstetrician/ gynecologists. *Obstet Gynecol* 2001;98:7-13.
- Wedgwood, J. F.; Carlin, E. B.; Benjamin, B. L.; Weintraub, A.; Lasker, M.; Green, R. S., and Holzman, I. R. Penicillin at birth can help prevent early-onset group B streptococcal disease. *Pediatrics.* 1997 Apr; 99(4):651-2.
- Wendel, G. D. Jr; Leveno, K. J.; Sanchez, P. J.; Jackson, G. L.; McIntire, D. D., and Siegel, J. D. Prevention of neonatal group B streptococcal disease: A combined intrapartum and neonatal protocol. *Am J Obstet Gynecol.* 2002 Apr; 186(4):618-26.
- Wiswell TE, Baumgart S, Gannon CM, Spitzer AR. No lumbar puncture in the evaluation for early neonatal sepsis: will meningitis be missed? *Pediatrics* 1995;95:803-6.

- Wiswell TE, Stoll BJ, Tuggle JM. Management of asymptomatic, term gestation neonates born to mothers treated with intrapartum antibiotics. *Pediatr Infect Dis J* 1990;9:826–31.
- Wood EG, Dillon HC. A prospective study of group B streptococcal bacteriuria in pregnancy. *Am J Obstet Gynecol* 1981;140:515–20.
- Yancey MK, Armer T, Clark P, Duff P. Assessment of rapid identification tests for genital carriage of group B streptococci. *Obstet Gynecol* 1992;80:1038–47.
- Yancey MK, Duff P, Clark P, Kurtzer T, Frentzen BH, Kubilis P. Peripartum infection associated with vaginal group B streptococcal colonization. *Obstet Gynecol* 1994;84:816–9.
- Yancey MK, Duff P, Kubilis P, Clark P, Frentzen BH. Risk factors for neonatal sepsis. *Obstet Gynecol* 1996;87:188–94.
- Yancey MK, Schuchat A, Brown LK, Ventura VL, Markenson GR. The accuracy of late antenatal screening cultures in predicting genital group B streptococcal colonization at delivery. *Obstet Gynecol* 1996;88:811–5.
- Zaleznik DF, Rench MA, Hillier S, et al. Invasive disease due to group B streptococcus in pregnant women and neonates from diverse population groups. *Clin Infect Dis* 2000;30:276–81.
- Zangwill KM, Schuchat A, Wenger JD. Group B streptococcal disease in the United States, 1990: report from a multistate active surveillance system. *MMWR* 1992;41(SS-6):25–32.